









Salinisation process of Lake Sidi Boughaba, Kenitra, North Western, Morocco: A statistical approach

Mohamed Lachhab¹ , Mohamed Najy¹ , Fatima Zahra Talbi^{2) 3)} , Aziz Taouraout¹ , Mohamed El Qryefy¹ , Hassan Ech-Chafay¹ , Driss Belghyti¹ , Khadija El Kharrim¹ 

¹⁾ University Ibn Tofail, Faculty of Sciences, Natural Resources and Sustainable Development Laboratory, BP 133, 14000 Kenitra, Morocco

²⁾ Hassan First University of Settat, Faculty of Sciences and Technologies, Laboratory of Biochemistry, Neurosciences, Natural Resources and Environment, Settat, Morocco

³⁾ Sidi Mohamed Ben Abdellah University, Faculty of Sciences Dhar El Mahraz, Laboratory of Biotechnology, Conservation and Valorization of Natural Resources (LBCVNR), Fez, Morocco

RECEIVED 21.11.2020

ACCEPTED 18.09.2021

AVAILABLE ONLINE 28.09.2022

Abstract: In order to determine the nature of the high salinisation rate of the waters of Lake Sidi Boughaba, which has been listed in the Ramsar list since 1980, 23 samples that were taken during four sampling operations were subjected to physicochemical analyses. The obtained results were processed using a combination of bi-varied methods (correlation tests) and multivariate statistical methods (principal component analysis – PCA). The physicochemical analyses reveal that they are alkaline waters with a pH ranging between 8.38 and 9.03, an electrical conductivity (EC) of the order of 12.4 to 17.4 mS·cm⁻¹, and high levels of Na⁺ and Cl⁻, up to 3700 and 6630 mg·dm⁻³ respectively, indicating a marine origin of these waters. In addition, the statistical treatment revealed that the mineralisation of the waters of this ecosystem is controlled by four main mechanisms of the salinisation; the main mechanism underlying this strong mineralisation is due to the impact of the marine spray. The second-order processes are about the phenomenon of the ion exchange, the dissolution/precipitation of evaporitic and carbonate formations, the oxidation–reduction processes, notably the reduction of sulphates as well as biochemical phenomena due to the selective absorption of certain ions by fauna and flora.

Keywords: Kenitra, marine aerosols, Morocco, salinisation, Sidi Boughaba Lake

INTRODUCTION

Salinisation is one of the leading causes of water quality degradation in the world [VENGOSH 2003]. In the marine domain, marine intrusion is one of the most common salinisation mechanisms affecting coastal water quality [JONES *et al.* 1999]. However, the sea spray that is a kind of aerosol removed by the wind at the crest of the waves, in turn, constitutes a potential natural source of salinisation for coastal waters [YOUNSI 2001]. In Morocco, the conditions of the arid climate and their hydrological consequences are the fundamental factors behind the formation of salty waters. The salinity of the waters of Sidi Boughaba Lake and the overlying groundwater have been the subject of a number of previous studies to explain the origin, causes, and the evolution of this phenomenon [BEN KABBOUR 2002; MARGAT 1961; RAMDANI 1981; ZOUHRI *et al.* 2010]. The classic approach to interpreting

hydrochemical variables is limited to analysing hydrochemical processes and classifying hydrochemical facies using various graphic methods. In contrast, multivariate statistical techniques are the best and, most often, the only effective solution for analysing a large body of information generated from hydrochemical data [JOIN *et al.* 1997]. In this respect, principal component analysis (PCA) was performed to better constrain the interpretation of data using the XLSTAT 2019 statistical analysis software.

MATERIALS AND METHODS

SITE DESCRIPTION

The Sidi Boughaba reserve occupies a coastal interdune basin of 6 km long and 300–800 m wide parallel to the Atlantic coast, over an area of 650 ha. It consists of a succession of dunes oriented

from N-NE to S-SW (Fig. 1) separated in the middle by a depression occupied by a lake [CHERKAOUI, BOUCHAFRA 2003]. The formation of this lacustrine environment dates back to 6520 ±110 before present [REILLE 1979]. This age corresponds to the Mellahian phase of the Moroccan Quaternary (Flanders of Europe) [ATBIB 1983]. The southern part is locally colonised by chenopods. The dune ridges bordering the water body are occupied by *Eucalyptus* sp., *Juniperus phoenicea*, *Pistacia lentiscus*, *Retama monosperma*, *Populus alba*, *Phillyrea angustifolia*, *Chamaerops humilis*, *Tamarix gallica* and *Olea europaea* are relatively undisturbed [RAMDANI et al. 2001]. The climate of this area is characterised by the influence of the Atlantic Ocean. The rainfall pattern is characterised by a seasonal distribution of precipitation. The annual average rainfall height is 557.2 mm. The average levels of temperature in August and January are approximately 23.8 and 12.2°C, respectively. Weak white frosts can occur in winter. The annual mean evapotranspiration value is 837.11 mm. It is greater than the precipitation value [BELGHAZI, MOUNIR 2016; NAJY et al. 2021]. The lake is fed mainly by the water table [CHERKAOUI, BOUCHAFRA 2003]. Its water is brackish with salinity ranging from 3 to 27 g·dm⁻³ depending on the season [LACHHAB et al. 2013].




Fig. 1. Descriptive map of the Lake Sidi Boughaba wetland; source: own elaboration

SAMPLING AND ANALYSIS METHODS

The data presented in this study come from four water sampling campaigns carried out on six stations from July, 2009 to May, 2010 (Tab. 1).

Table 1. Description of sampling stations

Station	Description
S1	located in the north of the permanent part
S2	located in the middle of the permanent part
S3	located in the south of the permanent part
S4	located in the northern part of the lake and separated from the permanent water body by a road that leads back to the center of education
S5	located at the south end of the lake in a semi-permanent biotope
S6	located outside the reserve in a temporary biotope

Source: own elaboration.

The samples were filtered through a 0.45 µm filter and were collected in polyethylene bottles with a capacity of 2 dm³ without air bubbles, and those for cation determination were acidified using nitric acid to prevent adsorption and precipitation of the chemical elements and were stored in portable coolers (+4°C) [RODIER 2009]. Then, they were analysed on their arrival at the National Institute of Hygiene's (INH) (Fr. Institut National d'Hygiène) hydrology laboratory in Rabat. The analyses of various physicochemical parameters were carried out according to the method described by RODIER [2009]. Reaction (pH), temperature (*T*), dissolved oxygen (*DO*), electrical conductivity (*EC*) and were total dissolved solids (*TDS*) measured *in situ* by the use of a portable pH meter, type (WTW pH 330i/SET), thermometer incorporated in the pH meter, Orion Model 607A Oximeter, Portable Conductivity Meter, type (WTWcond330i/SET) and salinometer incorporated in the conductivity meter. Furthermore, the major cations (Na⁺, Mg²⁺, Ca²⁺ and K⁺) were determined by the VARIAN atomic absorption spectrophotometer model, the two anions (Cl⁻ and SO₄²⁻) were measured by Dionex 2000 ionic chromatography, the biological oxygen demand (*BOD*₅) by BOD meter, the total alkalinity (*TA*) and bicarbonates (HCO₃⁻) with a volumetric dosing of 0.1 N HCl and colour by Nessleriser Comparator.

STATISTICAL METHODS OF DATA PROCESSING

The classical approach to the interpretation of hydrochemical variables considers hydrochemical variables as limited to interpreting hydrochemical processes and classifying hydrochemical facies using various graphical methods [ABOUBAKAR 2012]. Multivariate statistical techniques such as principal component analysis are better and often the only effective solution for analysing a large mass of information [JOIN et al. 1997].

Principal component analysis (PCA) is a mathematical procedure that uses an orthogonal transformation to convert a set of observations of possibly correlated variables into a set of uncorrelated variables called principal components [QIN et al. 2013]. It reduces a large number of variables to a small number of main components by linearly combining measurements made on the original variables [GULER et al. 2012]. The first step is to calculate the correlation coefficients between the variables. Subsequently, the projection axes are assimilated to the eigenvectors of the correlation matrix [BEN ALAYA et al. 2014]. The number of components to be extracted is determined by the variance accumulation test commonly called "Scree test", where the extraction of the components at the location of the slope change in the eigenvalue graph [Aw et al. 2011]. This statistical technique has been widely applied to investigate environmental phenomena and hydrogeochemical processes around the world [Aw et al. 2011; BEN ALAYA et al. 2014; EBLIN et al. 2014; MOUISSI, ALAYAT 2016; REGGAM et al. 2015] and in Morocco [EL YAOUTI et al. 2009; FADILI et al. 2015; NAJIB et al. 2016; 2017].

The PCA has 6 descriptors and 15 variables which are: *EC*, pH, *T*, *DO*, *TA*, colour, *BOD*₅, *TDS*, and ions such as HCO₃⁻, SO₄²⁻, Cl⁻, Na⁺, K⁺, Ca²⁺ and Mg²⁺. It was carried out by using the software XLSTAT 2019. The combination of all the methods used in this study makes it possible to know the origin of the salinisation process of the Sidi Boughaba Lake waters.

RESULTS AND DISCUSSION

DESCRIPTIVE STATISTICS

The results of physicochemical analyses carried out on the waters of Sidi Boughaba Lake (Tab. 2) reveal that the T varies between 19.46 and 21.55°C, with an average of 21.04 ±0.78°C. The pH values are ranging between 8.38 and 9.03, with an average of 8.72 ±0.3. The EC ranges from 12.4 to 17.4 mS·cm⁻¹, with an average value of 14.12 ±2.19 mS·cm⁻¹.

Table 2. Descriptive statistics of the physicochemical parameters of the waters of the Lake Sidi Boughaba

Parameter	Unit	Value			
		min.	max.	mean ±SD	CV
T	°C	19.46	21.55	21.04 ±0.78	0.04
DO	mg·dm ⁻³	4.81	8.58	6.86 ±1.28	0.19
pH	pH unit	8.38	9.03	8.72 ±0.30	0.03
EC	mS·cm ⁻¹	12.4	17.4	14.12 ±2.19	0.16
Colour	mg Pt·dm ⁻³	71.25	475.00	141.60 ±163.41	1.15
TDS	g·dm ⁻³	7.59	11.23	8.91 ±1.62	0.18
BOD_5	mg·dm ⁻³	5.00	7.50	6.21 ±1.05	0.17
TA	meq·dm ⁻³	5.12	18.29	8.29 ±4.95	0.60
Na^+	mg·dm ⁻³	2526	3700	3019.67 ±506.72	0.17
Ca^{2+}	mg·dm ⁻³	46.69	163.2	97.47 ±49.47	0.51
Mg^{2+}	mg·dm ⁻³	271.33	485.05	340.50 ±81.72	0.24
K^+	mg·dm ⁻³	38.64	57.66	45.72 ±7.27	0.16
Cl^-	mg·dm ⁻³	5007	6630	5595.00 ±678.12	0.12
SO_4^{2-}	mg·dm ⁻³	139.95	1091.07	502.79 ±397.60	0.79
HCO_3^-	mg·dm ⁻³	303.18	1001.96	484.86 ±257.82	0.53

Explanations: SD = standard deviation, CV = coefficient of variation, T = temperature, DO = dissolved oxygen, EC = electrical conductivity, TDS = total dissolved solids, BOD_5 = biological oxygen demand, TA = total alkalinity.

Source: own study.

DO levels range from 4.81 to 8.58 mg·dm⁻³, which is considered as an average of 6.86 ±1.28 mg·dm⁻³. The TDS rate varies between 7.59 and 11.23 g·dm⁻³, which is considered an average of 8.91 ±1.62 g·dm⁻³. The BOD_5 has a minimum of 5 mg·dm⁻³ and a maximum of 7.5 mg·dm⁻³, with an average value of 6.21 ±1.05 mg·dm⁻³. The TA oscillates between 5.12 and 18.29 meq·dm⁻³, which is considered an average of 8.29 ±4.95 meq·dm⁻³. The colour varies between 71.25 and 475 mg Pt·dm⁻³, with an average of 141.60 ±163.41 mg Pt·dm⁻³. Sodium and magnesium

appear to be the dominant cations ($Na^+ > Mg^{2+} > Ca^{2+} > K^+$), while chlorides and bicarbonates are the dominant anions ($Cl^- > HCO_3^- > SO_4^{2-}$). Concentration of Na^+ ranges from 2526 to 3700 mg·dm⁻³ with an average of 3019.67 mg·dm⁻³. Mg^{2+} levels range from a high rate of 485.05 mg·dm⁻³ to a low rate of 271.33 mg·dm⁻³ with an average of 340.50 mg·dm⁻³. Ca^{2+} ions have a maximum of 163.2 mg·dm⁻³ and a minimum of 46.69 mg·dm⁻³ with an average of 97.47 whereas K^+ values range from 38.64 mg·dm⁻³ to 57.66 mg·dm⁻³ with an average value of 45.72 mg·dm⁻³. In addition, Cl^- concentrations range from 5007 to 6630 mg·dm⁻³ with an average of 5595 mg·dm⁻³; HCO_3^- concentrations vary between a maximum of 1001.96 mg·dm⁻³ and a minimum of 303.18 mg·dm⁻³ with an average value of 484.86 mg·dm⁻³ while the SO_4^{2-} values are between 139.95 mg·dm⁻³ and 1091.07 mg·dm⁻³.

CORRELATION BETWEEN VARIABLES


The study of bi-variate linear correlations between the studied physicochemical parameters proves the solid associations between them. The different relations between the parameters controlled by the estimation of the correlation coefficient can be determined on the basis of the theoretical critical coefficient drawn from the table of Bravais–Pearson according to the number of couples put into play. On this basis, the critical coefficient is 0.4134 for the number of pairs 21, and the values, the absolute values of which are greater than said critical coefficient, are written in bold (Tab. 3).

The T shows a slight positive correlation with the BOD_5 of 0.596, which could be explained by a decomposition of organic matter [NÉGREL *et al.* 2001]. It also shows a negative correlation successively with Ca^{2+} (-0.672) and DO (-0.634), this can be explained by a strong precipitation of calcium carbonates under the impact of evaporation and the decrease of the oxygen solubility. DO is positively correlated with pH (0.437) and Ca^{2+} (0.468). It also has a negative correlation with TDS , Mg^{2+} , Na^+ , HCO_3^- , Cl^- , colour, TA , and BOD_5 .

The correlation between pH and DO is one of the essential characteristics of eutrophic environments both for phytoplankton development and for macrophytes [Aw *et al.* 2011]. The colour is positively correlated with TA , HCO_3^- , SO_4^{2-} , Na^+ and Cl^- as well as a negative correlation with Ca^{2+} . BOD_5 is positively correlated with TA and HCO_3^- and negatively with Ca^{2+} .

Strong correlations have been observed between TDS and EC with a value of Na^+ , Mg^{2+} , K^+ , Cl^- , SO_4^{2-} and HCO_3^- . This strong correlation implies that these ions participated in the mineralisation of the waters of this lacustrine environment [ASKRI *et al.* 2016]. They also demonstrate a slight positive correlation with colour.

HCO_3^- ions are positively correlated with Na^+ and SO_4^{2-} . The association with Na^+ could be explained by chemical reactions and mineral dissolution [GANYAGLO *et al.* 2010], whereas the correlation with SO_4^{2-} could be explained by the reduction of SO_4^{2-} ions to HCO_3^- , according to the following reaction [DE MONTEY *et al.* 2008]:



They also have a negative correlation with calcium. This can be explained by the complexation of Ca– HCO_3 and saturation with

Table 3. Correlation matrix for all data ($n = 23$)

Parameter	<i>T</i>	<i>DO</i>	<i>pH</i>	<i>EC</i>	<i>Colour</i>	<i>TDS</i>	<i>BOD₅</i>	<i>TA</i>	Na^+	Ca^{2+}	Mg^{2+}	K^+	Cl^-	SO_4^{2-}	HCO_3^-
<i>T</i>	1														
<i>DO</i>	-0.634	1													
<i>pH</i>	0.372	0.437	1												
<i>EC</i>	-0,008	-0.611	-0.892	1											
<i>Colour</i>	0.168	-0.776	-0.586	0.497	1										
<i>TDS</i>	-0,082	-0.581	-0.927	0.996	0.514	1									
<i>BOD₅</i>	0.596	-0.676	0.061	-0.005	0.584	-0.022	1								
<i>TA</i>	0.320	-0.818	-0.448	0.403	0.984	0.408	0.683	1							
Na^+	0.057	-0.705	-0.882	0.974	0.581	0.972	0.198	0.501	1						
Ca^{2+}	-0.672	0.468	-0.423	0.368	-0.479	0.393	-0.768	-0.619	0.252	1					
Mg^{2+}	-0,257	-0.271	-0.832	0.909	0.132	0.913	-0.337	0.007	0.840	0.715	1				
K^+	-0,374	-0.198	-0.855	0.877	0.141	0.892	-0.293	0.002	0.830	0.742	0.976	1			
Cl^-	0.296	-0.687	-0.670	0.922	0.458	0.890	0.081	0.415	0.882	0.165	0.766	0.702	1		
SO_4^{2-}	-0,371	-0.465	-0.961	0.792	0.744	0.834	0.039	0.616	0.794	0.226	0.657	0.691	0.580	1	
HCO_3^-	0.354	-0.827	-0.419	0.386	0.977	0.390	0.700	0.999	0.487	-0.644	-0.015	-0.022	0.411	0.588	1

Explanations: *T*, *DO*, *EC*, *TDS*, *BOD₅*, *TA* as in Tab. 2.
Source: own study.

CaCO_3 [KETATA *et al.* 2011]. Chlorides show a strong positive correlation with the major components of seawater (Na^+ and SO_4^{2-}) whereas $\text{Cl}-\text{Na}$ ($r = 0.882$) and $\text{Cl}-\text{SO}_4$ ($r = 0.580$) indicate that seawater influenced the lake water mineralisation [ASKRI *et al.* 2016]. The above-mentioned significant correlation between Na^+ and Cl^- as well as the association between Mg^{2+} and Cl^- ($r = 0.766$) takes place due to seawater intrusion [MTONI *et al.* 2013]. The strong correlation between K^+ and Cl^- ($r = 0.702$) and the moderate correlation between K^+ and SO_4^{2-} ($r = 0.691$) could be explained by marine intrusion [ASKRI *et al.* 2016; MTONI *et al.* 2013; NAJIB *et al.* 2017]. The low correlation observed between Ca^{2+} and Cl^- results from the removal of calcium following precipitation of calcite [MTONI *et al.* 2013]. The strong correlation between Mg^{2+} and Ca^{2+} ($r = 0.715$) can be attributed to the dissolution of calcite and dolomite minerals [ASKRI *et al.* 2016; KETATA *et al.* 2011; NAJIB *et al.* 2017; QIN *et al.* 2013] and the exchange of cations resulting from the intrusion of seawater [NAJIB *et al.* 2017; QIN *et al.* 2013]. The ions are strongly correlated with Mg^{2+} ($r = 0.840$). This result would be attributed to ion exchange reactions [ASKRI *et al.* 2016]. The correlation of potassium with sodium, magnesium, and calcium is explained by the phenomenon of inverse ionic exchange between water and aquifer and which is resulting from the adsorption of Na^+ or K^+ and the release of Ca^{2+} or Mg^{2+} [ASKRI *et al.* 2016; CAPACCIONI *et al.* 2005; IBRAHIM *et al.* 2018; NAJIB *et al.* 2016;]. Sulphate ions are correlated with sodium and magnesium. The lack of correlation between Ca^{2+} and SO_4^{2-} seeks an origin for calcium other than a dissolution of gypsum [ABOUBAKAR 2012; GANYAGLO *et al.* 2010].

PRINCIPAL COMPONENT ANALYSIS (PCA)

The results obtained with principal component analysis (PCA) are reported in Tables 4 and 5. Based on the variance accumulation test commonly known as Scree test [Aw *et al.* 2011], two main factors were extracted, the sum of the variances

Table 4. Neat values and percentages expressed by the main axes

Parameter	Value for factor				
	F1	F2	F3	F4	F5
Eigenvalue	8.244	4.899	1.427	0.349	0.081
Variability (%)	54.962	32.663	9.512	2.324	0.539
Cumulative (%)	54.962	87.625	97.137	99.461	100.000

Source: own study.

Table 5. The contribution of variables in the determination of factors

Parameter	Value for factor				
	F1	F2	F3	F4	F5
<i>T</i>	0.007	-0.685	0.721	-0.089	-0.056
<i>DO</i>	-0.721	0.612	-0.281	-0.020	0.164
<i>pH</i>	-0.922	-0.266	0.271	-0.042	0.065
<i>EC</i>	0.961	0.209	0.175	-0.041	0.011
<i>Colour</i>	0.710	-0.603	-0.355	-0.080	0.021
<i>TDS</i>	0.967	0.233	0.096	-0.022	0.001
<i>BOD₅</i>	0.186	-0.855	0.077	0.473	0.073
<i>TA</i>	0.623	-0.731	-0.266	-0.082	0.014
Na^+	0.975	0.073	0.156	0.142	-0.004
Ca^{2+}	0.172	0.981	0.006	0.086	-0.032
Mg^{2+}	0.784	0.586	0.191	0.018	-0.065
K^+	0.771	0.609	0.077	0.155	0.060
Cl^-	0.863	0.037	0.428	-0.198	0.176
SO_4^{2-}	0.887	0.081	-0.452	-0.043	-0.019
HCO_3^-	0.606	-0.754	-0.240	-0.082	0.013

Explanations: *T*, *DO*, *EC*, *TDS*, *BOD₅*, *TA* as in Tab. 2.
Source: own study.

expressed is 87.625% (Tab. 4). These factors truly reflect the variance expressed and explain the information sought. The representation using these two factors accounts for the structure of the scatter plots satisfactorily. The results of the PCA make it possible to choose the different necessary elements for the interpretation of the different data. The analysis of the space graphs of the variables in the factorial plane F1–F2 (Fig. 2) shows that this plane alone represents 87.62% of the researched information as well as the expressed variance. The circle of community (Fig. 2) shows that the factor F1, the most important, is positively correlated with *EC*, *TDS*, Mg^{2+} , Na^+ , K^+ , Cl^- and SO_4^{2-} (Tab. 5), which can also be attributed to the intrusion of sea water [ASKRI *et al.* 2016]. Elevated levels of *TDS*, Cl^- and Na^+ are generally from marine sources, either seawater intrusion into aquifers or marine aerosols carried by precipitation [QIN *et al.* 2013]. It is also positively correlated with colour, *TA*, and HCO_3^- ions as well as a negative correlation with *DO* and *pH* (Tab. 5). Factor F2 shows a negative correlation with *T*, *pH*, colour, *BOD*₅, *TA*, and HCO_3^- ions, as opposed to *DO* and cations Ca^{2+} , Mg^{2+} and K^+ , which could take place due to a rise in temperature which is responsible for the precipitation of calcite; this precipitation has an influence on a variety of chemical parameters: bicarbonates, dissolved CO_2 , Ca^{2+} , and consequently salinity, conductivity, alkalimetric title (*AT*), complete alkalimetric title (*CAT*), hydrotimetric title (*HT*) and hardness [ROCHE 1979]. The opposite evolution of Ca^{2+} and HCO_3^- indicates another source in addition to carbonate alteration, such as gypsum dissolution and cation exchange, which increases the Ca^{2+} content [EL YAOUTI *et al.* 2009]. The positive correlation between the three cations (Mg^{2+} , Ca^{2+} , and K^+) could be explained by the absorption of these elements by the plants during this period which knows a great photosynthetic activity. Seasonal growth of molluscs, which is faster in warmer waters, is expected to result in higher calcium absorption from April to October, especially during the low water level period [ROCHE 1979]. It may also take place due to the uptake of these elements by vegetation [ROCHE 1979]. The graphical representation in the factorial space of the statistical units (Fig. 3) is due to the distribution of the water points




Fig. 2. Space of the variables in the factorial plane F1–F2; source: own study




Fig. 3. Space of individuals in the factorial plane F1–F2; source: own study

according to the different factors F1–F2. The analysis of this graph shows a single grouping of the water points, which reflects the homogeneity of the physicochemical characteristics of the waters of this lacustrine environment.

CONCLUSIONS

The statistical study carried out on the hydrochemical variables resulting from the analysis of the waters of the Lake Sidi Boughaba made it possible to classify this lake environment as a eutrophic medium. It also revealed high levels of *TDS*, Cl^- and Na^+ on the one hand, and their correlation indicating a marine source, be it the intrusion of seawater into the aquifers or the marine aerosols carried by the precipitations on the other hand. It identified four main mechanisms that are responsible for the evolution of the mineralisation of the waters in the area. The mechanism at the base of the mineralisation in major elements of the waters is the effect of sea spray. The second-order processes are the ion exchange phenomenon, the dissolution/precipitation of evaporitic and carbonate formations, the oxidation-reduction processes, notably the reduction of sulphates as well as biochemical phenomena due to the selective absorption of certain ions by fauna and flora.

ACKNOWLEDGMENTS

The authors would like to thank Miss Soukaina AROUB for the assistance in the field of computing and Mr. Miloud CHAKIT and Mr. Mohamed HA-NINE for the assistance in the field of sampling.

REFERENCES

- ABOUBAKAR M. 2012. Caractérisation d'un système aquifère volcanique par approche couplée hydrogéochimique et modélisation numérique. Exemple de l'aquifère des basaltes de Dalha, sud-ouest de

- la République de Djibouti [Characterisation of a volcanic aquifer system by coupled hydrogeochemical approach and numerical modeling. Example of the basalt aquifer of Dalha, southwest of the Republic of Djibouti]. PhD Thesis. Poitiers. University of Poitiers pp. 238.
- ASKRI B., AHMED A.T., AL-SHANFARI R.A., BOUHLILA R., AL-FARISI K.B. 2016. Isotopic and geochemical identifications of groundwater salinisation processes in Salalah coastal plain, Sultanate of Oman. *Chemie der Erde – Geochemistry*. Vol. 76. No. 2. DOI 10.1016/j.chemer.2015.12.002.
- ATBIB M. 1983. Etude phytocécologique de la réserve biologique de Mehdia (littoral Atlantique du Maroc). 2 – La végétation du milieu dunaire [Phytoecological study of the biological reserve of Mehdia (Atlantic coast of Morocco). 2 – The vegetation of the dune environment]. *Bulletin de l'Institut Scientifique. Supplement 7. Université Mohammed V. Institut Scientifique. Rabat* pp. 112.
- AW S., N'GORAN E.B.Z., SIAKA S., PARINET B. 2011. Intérêt de l'analyse multidimensionnelle pour l'évaluation de la qualité physico-chimique de l'eau d'un système lacustre tropical: cas des lacs de Yamoussoukro (Côte d'Ivoire) [Interest of multidimensional analysis for the assessment of the physico-chemical quality of water in a tropical lake system: Case of the Yamoussoukro lakes (Ivory Coast)]. *Journal of Applied Biosciences*. Vol. 38 p. 2573–2585.
- BELGHAZI B., MOUNIR F. 2016. Analyse de vulnérabilité au changement climatique du couvert forestier, Forêt de la Maâmora (Maroc) [Analysis of vulnerability to climate change of the forest cover, Maâmora Forest (Morocco)]. *Rapport technique. Rome. FAO. ISBN 978-92-5-109471-6* pp. 124.
- BEN ALAYA M., ZEMNI T., MAMOU A., ZARGOUNI F. 2014. Acquisition de salinité et qualité des eaux d'une nappe profonde en Tunisie: approche statistique et géochimique [Acquisition of salinity and water quality from a deep aquifer in Tunisia: statistical and geochemical approach]. *Journal des Sciences Hydrologiques*. Vol. 59(2) p. 395–419. DOI 10.1080/02626667.2013.870663.
- BEN KABBOUR B. 2002. Exploration, évaluation et protection des eaux souterraines en zones côtières marocaines: approches géophysique, hydrogéochimique, modélisation et S.I.G., cas de la Mamora occidentale [Exploration, evaluation and protection of groundwater in Moroccan coastal areas: geophysical, hydrogeochemical approaches, modeling and S.I.G., case of the western Mamora]. PhD Thesis. Kenitra. Ibn Tofail University pp. 300.
- CAPACCIONI B., DIDERO M., PALETTA C., DIDERO L. 2005. Saline intrusion and refreshing in a multilayer coastal aquifer in the Catania plain (Sicily, southern Italy): dynamics of degradation processes according to the hydrochemical characteristics of groundwaters. *Journal of Hydrology*. Vol. 307 p. 1–16.
- CHERKAoui I., BOUCHAFRA A. 2003. Fiche descriptive sur les zones humides Ramsar (FDR) [Ramsar wetlands fact sheet (RIS)]. Mehdia. National Center for Environmental Education (Morocco) [online]. [Access 30.08.2020]. Available at: <https://rsis.ramsar.org/RISapp/files/RISrep/MA207RIS.pdf>
- DE MONTETY V., RADAKOVITCH O., VALLET-COULOMB C., BLAVOUX B., HERMITTE D., VALLES V. 2008. Origin of groundwater salinity and hydrogeochemical processes in a confined coastal aquifer: Case of the Rhône delta (Southern France). *Applied Geochemistry*. Vol. 23 p. 2337–2349.
- EBLIN S.G., SOMBO A.P., SORO G., AKA N., KAMBIRÉ O., SORO N. 2014. Hydrochimie des eaux de surface de la région d'Adiaké (sud-est côtier de la Côte d'Ivoire) [Hydrochemistry of surface water in the Adiaké region (south-eastern coast of Côte d'Ivoire)]. *Journal of Applied Biosciences*. Vol. 75 p. 6259–6271. DOI 10.4314/jab.v75i1.10.
- EL YAOUTI F., EL MANDOUR A., KHATTACH D., BENAVENTE J., KAUFMANN O. 2009. Salinization processes in the unconfined aquifer of Bou-Areg (NE Morocco): A geostatistical, geochemical, and tomographic study. *Applied Geochemistry*. Vol. 24 p. 16–31.
- FADILI A., MEHDI K., RISS J., NAJIB S., MAKAN A., BOUTAYAB K. 2015. Evaluation of groundwater mineralization processes and seawater intrusion extension in the coastal aquifer of Oualidia, Morocco: Hydrochemical and geophysical approach. *Arabian Journal of Geosciences*. DOI 10.1007/s12517-015-1808-5.
- GANYAGLO S.Y., BANOENG-YAKUBO B., OSAE S., DAMPARE S.B., FIANKO J.R. 2010. Water quality assessment of groundwater in some rock types in parts of the eastern region of Ghana. *Environmental Earth Sciences*. Vol. 62 p. 1055–1069.
- GULER C., ALI KURT M., ALPASLAN M., AKBULUT C. 2012. Assessment of the impact of anthropogenic activities on the groundwater hydrology and chemistry in Tarsus coastal plain (Mersin, SE Turkey) using fuzzy clustering, multivariate statistics and GIS techniques. *Journal of Hydrology*. Vol. 414–415 p. 435–451.
- IBRAHIM R.G.M., KORANY E.A., TEMPEL R.N., GOMAA M.A. 2018. Processes of water-rock interactions and their impacts upon the groundwater composition in Assiut area, Egypt: Applications of hydrogeochemical and multivariate analysis. *Journal of African Earth Sciences*. Vol. 149. DOI 10.1016/j.jafrearsci.2018.07.026.
- JOIN J., COUDRAY J., LONGWORTH K. 1997. Using principal components analysis and Na/Cl ratios to trace groundwater circulation in a volcanic island: the example of Reunion. *Journal of Hydrology*. Vol. 190(2) p. 1–18.
- JONES B.F., VENGOSH A., ROSENTHAL E., YECHIELI Y. 1999. Géochimical investigations. In: *Seawater intrusion in coastal aquifers – concepts, methods and practices*. Eds. J. Bear, A. H.-D. Cheng, S. Sorek, D. Ouazar, I. Herrera. Dordrecht. Springer Science +Business Media p. 51–72.
- KETATA M., HAMZAOUIA F., GUEDDARI M., BOUHLILA R., RIBEIRO L. 2011. Hydrochemical and statistical study of groundwaters in Gabes-South deep aquifer (south-eastern Tunisia). *Physics and Chemistry of the Earth*. Vol. 36 p. 187–196.
- LACHHAB M., BELGHYTI D., NAJY M., FATIMETOU SALMA M.M.O.S., ECH-CHAFAÏ H., AKKAoui O., BEN AAKAME R., EL KHARRIM K. 2017. A coastal lake's hydro-chemistry: Sidi Boughaba Case, North Western Morocco. *International Journal of Scientific & Engineering Research*. Vol. 8(3) p. 1904–1909.
- LACHHAB M., EL KHARRIM K., EL ABIDI A., BEN AKKAME R., BELGHYTI D. 2013. Etude physico-chimique des eaux du lac Sidi Boughaba – Site Ramsar – Kénitra Maroc [Physico-chemical study of the waters of Lake Sidi Boughaba – Ramsar site – Kenitra Morocco]. *ScienceLib Editions Mersenne*. Vol. 5. No. 131216 p. 1–13.
- MARGAT J. 1961. Les eaux salées au Maroc: Hydrogéologie et hydrochimie [Salt water in Morocco: Hydrogeology and hydro-chemistry]. *Notes et Mémoires. Service géologique du Maroc*. No. 151 pp. 137.
- MOUISSI S., ALAYAT H. 2016. Utilisation de l'analyse en composantes principales (ACP) pour la caractérisation physico-chimique des eaux d'un écosystème aquatique: Cas du lac Oubéira (Extrême NE Algérie) [Use of principal component analysis (PCA) for the physico-chemical characterisation of the waters of an aquatic ecosystem: The case of Lake Oubéira (Extreme NE of Algeria)]. *Journal of Materials and Environmental Science*. Vol. 7(6) p. 2214–2220.
- MTONI Y., MJEMAH I.C., BAKUNDUKIZE C., VAN CAMP M., MARTENS K., WALRAEVENS K. 2013. Saltwater intrusion and nitrate pollution in

- the coastal aquifer of Dar es Salaam, Tanzania. *Environmental Earth Sciences*. Vol. 70(3) p. 1091–1111.
- NAJIB S., FADILI A., MEHDI K., RISS J., MAKAN A. 2017. Contribution of hydrochemical and geoelectrical approaches to investigate salinization process and seawater intrusion in the coastal aquifers of Chaouia, Morocco. *Journal of Contaminant Hydrology*. Vol. 198 p. 24–36. DOI 10.1016/j.jconhyd.2017.01.003.
- NAJIB S., FADILI A., MEHDI K., RISS J., MAKAN A., GUESSIR H. 2016. Salinization process and coastal groundwater quality in Chaouia, Morocco. *Journal of African Earth Sciences*. DOI 10.1016/j.jafrearsci.2015.12.010.
- NAJY M., LACHHAB M., ECH-CHAFAY H., TALBI F.Z., ACHHAR A., BELGHYTI D. 2021. A study on the salinization of coastal water of Sidi Boughaba Lake (Ramsar site 1980) in Morocco using the characteristic reports. *E3S Web of Conferences*. Vol. 234. DOI 10.1051/e3sconf/202123400016.
- NÉGREL P., BARBIER J., PETELET-GIRAUD E. 2001. Caractérisation géochimique et isotopique des relations eaux de surface – eaux souterraines de la zone humide de la Loire [Geochemical and isotopic characterisation of surface water – groundwater relationships in the Loire wetland]. *La Loire, Rapport BRGM. RP50668-FR* pp. 71.
- QIN R., WU Y., XU Z., XIE D., ZHANG C. 2013. Assessing the impact of natural and anthropogenic activities on groundwater quality in coastal alluvial aquifers of the lower Liaohe River Plain, NE China. *Applied Geochemistry*. Vol. 31 p. 142–158.
- RAMDANI M. 1981. Recherches hydrobiologiques sur la merja Sidi Boughaba, littoral atlantique du Maroc: étude physicochimique et analyse faunistique [Hydrobiological research on the merja Sidi Boughaba, Atlantic coast of Morocco: Physicochemical study and faunistic analysis]. *Bulletin de l'Institut Scientifique Rabat*. Vol. 5 p. 37–137.
- RAMDANI M., EL KHIATI N., FLOWER R.I., KRAIEM M.M., FATHI A.A., BIRKS H.H., PATRICK S.T. 2001. Open water zooplankton communities in North African wetland lakes: the CASSARINA Project. *Aquatic Ecology*. Vol. 35(3–4) p. 319–333. DOI 10.1023/A:1011957324901.
- REGGAM A., BOUCHELAGHEM H., HOUHAMDI M. 2015. Qualité physico-chimique des eaux de l'oued Seybouse (nord-est de l'Algérie): caractérisation et analyse en composantes principales [Physico-chemical quality of waters of Oued Seybouse (north-east of Algeria): Characterisation and analysis of principal components]. *Journal of Materials and Environmental Science*. Vol. 6(5) p. 1417–1425.
- REILLE M. 1979. Analyse pollinique du lac de Sidi Bou Rhaba, littoral Atlantique (Maroc) [Pollen analysis of Lake Sidi Bou Rhaba, atlantic coast (Morocco)]. *Ecologia Mediterranea*. Vol. 4 p. 61–65.
- ROCHE M.A. 1979. Traçage naturel salin et isotopique des eaux du système hydrologique du lac Tchad [Natural saline and isotopic tracing of the waters of the hydrological system of Lake Chad]. *Travaux et documents de L'O.R.S.T.O.M. No. 117*. ISBN 2-7099-0546-9 pp. 383.
- RODIER J. 2009. L'analyse de l'eau: eaux naturelles, eaux résiduaires et eaux de mer [Water analysis: Natural water, waste water and sea water]. 9th ed. Paris, Dunod. ISBN 978-2-10-054179-9 pp. 1526.
- VENGOSH A. 2003. Salinization and saline environments. In: *Environmental geochemistry. Treatise in geochemistry*. Eds. B. Lollar, H.D. Holland, K.T. Turekian. Elsevier Science p. 333–365.
- YOUNSI A. 2001. Méthodologie de mise en évidence des mécanismes de salure des eaux souterraines côtières en zone semi-aride irriguée (Chaouia Côtière, Maroc) [Methodology for demonstrating the salting mechanisms of coastal groundwater in irrigated semi-arid zone (coastal Chaouia, Morocco)]. PhD Thesis. El Jadida Université Chouaib Doukkali pp. 175.
- ZOUHRI L., TOTO E., CARLIER E., DEBIECHE T.H. 2010. Salinité des ressources en eau: intrusion marine et interaction eaux-roches (Maroc occidental) [Salinity of water resources: Marine intrusion and water-rock interaction (western Morocco)]. *Journal des Sciences Hydrologiques*. Vol. 55(8) p. 1337–1347.