










Plant secondary metabolites as defenses: A review

Herlina Jusuf¹⁾ , Marischa Elveny²⁾ , Feruza Azizova³⁾ ,
Rustem A. Shichiyakh⁴⁾ , Dmitriy Kulikov⁵⁾ , Muataz M. Al-Taee⁶⁾ ,
Karrar K. Atiyah⁷⁾ , Abduladheem T. Jalil⁸⁾ , Surendar Aravindhan⁹⁾ 

¹⁾ Universitas Negeri Gorontalo, Faculty of Sports and Health, Department of Public Health,
Jln. Jenderal Sudirman 6, Gorontalo, 96128, Indonesia

²⁾ Universitas Sumatera Utara, DS & CI Research Group, Medan, Indonesia

³⁾ Tashkent Medical Academy, Tashkent, Uzbekistan

⁴⁾ Kuban State Agrarian University named after I.T. Trubilin, Department of Management, Kuban, Russia

⁵⁾ Moscow State University of Technologies and Management named after K.G. Razumovsky (First Cossack University),
Department of Digital Nutrition, Hotel and Restaurant Services, Moscow, Russia

⁶⁾ AL-Nisour University College, Department of Medical Laboratories Technology, Baghdad, Iraq

⁷⁾ College of Dentistry, Al-Ayen University, Thi-Qar, Iraq

⁸⁾ Al-Mustaqbal University College, Medical Laboratories Techniques Department, Babylon, Hilla, Iraq

⁹⁾ Saveetha Institute of Medical and Technical Sciences, Chennai, India

RECEIVED 01.12.2021

ACCEPTED 08.03.2022

AVAILABLE ONLINE 31.12.2022

Abstract: Plant secondary metabolites have a variety of functions, including mediating relationships between organisms, responding to environmental challenges, and protecting plants against infections, pests, and herbivores. In a similar way, through controlling plant metabolism, plant microbiomes take part in many of the aforementioned processes indirectly or directly. Researchers have discovered that plants may affect their microbiome by secreting a variety of metabolites, and that the microbiome could likewise affect the metabolome of the host plant. Pesticides are agrochemicals that are employed to safeguard humans and plants from numerous illnesses in urban green zones, public health initiatives, and agricultural fields. The careless use of chemical pesticides is destroying our ecology. As a result, it is necessary to investigate environmentally benign alternatives to pathogen management, such as plant-based metabolites. According to literature, plant metabolites have been shown to have the ability to battle plant pathogens. Phenolics, flavonoids, and alkaloids are a few of the secondary metabolites of plants that have been covered in this study.

Keywords: alkaloids, flavonoids, phenolics, plant pathogens, secondary metabolites

INTRODUCTION

Since natural systems present plants with a variety of conflicting forces, plants have developed an extensive defensive system to cope with abiotic and biotic stresses [UL HAQ *et al.* 2019]. Because a multitude of stress factors interact to impact plants, every alteration in plant metabolic physiology cannot be attributed to a single stress component [GAŁCZYŃSKA *et al.* 2019; HEDJAL *et al.*

2018]. Various response pathways are activated in response to a particular stress, and there are several interconnections between signaling routes for herbivorous and pathogens insects [BODENHAUSEN, REYMOND 2007]. Some of these response mechanisms are activated independently of the antimicrobial nature, whereas others are triggered by infection [MANNIELLO *et al.* 2021]. Other plant defensive mechanisms include the creation of polymeric barriers to prevent pathogen entry and the production of

pathogen cell wall destroying enzymes (PCWDEs) [BELAIR *et al.* 2022; TYSKIEWICZ *et al.* 2022]. Plant pathogens generate PCWDEs as a subgroup of carbohydrate-active enzymes to deteriorate plant cell walls. Plants release PCWDE inhibitor proteins (PIPs) to limit the effect of PCWDEs [PRASANTH *et al.* 2022]. Plants also have unique signaling and identification mechanisms that allow them to quickly identify pathogen entry and mount an appropriate defensive response [WAHEED *et al.* 2022]. In the event that they become infected, plants have developed to adapt to further microbial attacks [TSUDA, KATAGIRI 2010; WU *et al.* 2014]. Secondary metabolism, which is distinct from primary metabolism, describes metabolic pathways and the tiny molecules they produce as byproducts that are not necessary for the growth and reproduction of the organism [DIXON *et al.* 2010]. Plant secondary metabolites (PSMs) are a diverse group of substances that are produced by secondary metabolic pathways in plants [DIVEKAR *et al.* 2022]. PSMs contain a significant number of structurally varied chemicals that were either produced as intermediates in the biosynthetic pathways of these primary metabolites or as primary metabolites themselves [JAMLOKI *et al.* 2021; YESHI *et al.* 2022]. A wide range of secondary metabolites are produced by plants that shield them from microbes and predators, as well as deter herbivores and microbes, depending on their poisonous nature. Certain secondary metabolites aid plant communication with other organisms, while others defend plants against abiotic stress such as metal toxicity, ozone, and ultraviolet (UV)-B, making secondary metabolites crucial for development and growth [CHAKRABORTY, ACHARYA 2017; KOZA *et al.* 2022]. CO₂ fixation enzymes, nitrogen metabolism, motility, pigmentation, survival, and growth are all known to be harmed by UV-B [PANDEY *et al.* 2020].

CLASSIFICATION

PSMs are typically categorized into six broad molecular families, depending on their biosynthesis pathways: phenolics, terpenes, steroids, alkaloids, and flavonoids [ZHANG *et al.* 2021]. Sulphur/

nitrogen (S/N), terpenes, and phenolics-containing compounds are the principal categories of secondary metabolites generated by plants (Fig. 1).

The greatest class of PSMs are terpenes. Some terpenes, such as gibberellins and brassinosteroids, are engaged in basic processes, such as plant growth and development, even though the majority of terpenes are crucial for plant defense [NINKUU *et al.* 2021; SHARMA *et al.* 2022]. A total of 25,000 different compounds are classified as terpenes, and they serve a variety of purposes such as preventing oviposition, feeding, or direct toxicity [ANSARI, AKHTAR 2019]. Terpenes have the ability to draw pollination insects. They have a fundamental unit called 5-C isoprene (isoterpenes, C₅H₈), which is toxic to herbivores [GALLI *et al.* 2005; HILLIER, LATHE 2019].

Nearly 10,000 different secondary metabolites make up the diverse category of chemicals known as plant phenols [BARTWAL *et al.* 2013]. Secondary metabolites with a hydroxyl functional group (phenyl group) attached to an aromatic ring are known as phenolics. They are the most widely dispersed secondary metabolites [VERMA, SHUKLA 2015]. Simple phenols (such as catechols and derivatives of hydroxybenzoic acid), lignins, condensed tannins, stilbenes, catechol melanins, and flavonoids are all examples of phenolic compounds [CHEYNIER *et al.* 2013]. Phenolics are a diverse group of molecules due to their unique chemical structures. In addition to aggressively deterring herbivores, these metabolites also actively entice pollinators to plants [BADRI *et al.* 2013]. Herbivores can be directly poisoned by phenols, or phenols can be oxidized by enzymes like polyphenol oxidases or peroxidases to produce toxic compounds that interfere physiologically with insect development and growth [HAMMER, BOWERS 2015].

Glucosides, which are mostly found in the Brassicaceae and Capparales plant taxa, are sulfur-containing secondary metabolites [WAR *et al.* 2020]. About 120 structural structures of the derivatives of amino acids known as glucosides were identified. The kind of glucoside is often determined by the side chain's amino acid precursor [BLAŽEVIĆ *et al.* 2020]. Younger leaves and plant reproductive parts have been found to have higher levels of




Fig. 1. Types of secondary metabolites; source: GARCIA-MIER *et al.* [2019]

glucosinolate [TSUNODA *et al.* 2017]. Usually, glucosinolates are found in the vacuoles of cells, where myrosinases have a protective coating over them (thioglucosidase). Damage caused by herbivores leads to the breakdown of plant cells, which in turn causes myrosinases to convert glucosinolates into poisonous compounds such as isothiocyanates, thiocyanates, and nitriles. These glucosinolate breakdown products have been demonstrated to be just as efficient as synthetic insecticides at deterring herbivorous insects from feeding by being extremely poisonous to them [KEITH, MITCHELL-OLDS 2017].

Alkaloids are among the second-generation metabolites that contain nitrogen [MICHAEL 2016]. The plant kingdom has reported about 10,000 distinct alkaloid compounds to date. On the basis of biosynthesis, alkaloids are separated into three classes: (a) proto-alkaloids, (b) pseudo-alkaloids, and (c) true alkaloids [KILGORE, KUTCHAN 2016]. Proto-alkaloids and real alkaloids both originate from amino acids. Pseudo-alkaloids, however, are not made from amino acids [LICHMAN 2021; RAJPUT *et al.* 2022]. True alkaloids share a heterocyclic ring with nitrogen and are derived from amino acids. Because alkaloids interfere with DNA replication, enzyme activity, and protein synthesis, they are poisonous to herbivores and disrupt neural signal transduction [BRIBI 2018].

Amino acids are the primary sources of sulphur and nitrogen-containing compounds [RODRIGUES-CORRÊA, FETT-NETO 2019]. *In vitro* studies of plants whose secondary metabolite expression was altered using current methods have demonstrated the defensive significance of secondary metabolites [LI *et al.* 2020]. The result of plant pathogen interactions over millions of years are secondary metabolites, and more than one hundred thousand metabolites are thought to be involved in the plant protection mechanism; thus, the situation is still unclear. Whereas large concentrations of secondary metabolites are expected to make plants better able to withstand biotic and abiotic stressors, for plant growth and reproduction, their synthesis is known to be costly [PIASECKA *et al.* 2015]. Numerous investigations revealed that hundreds of plant chemicals have chemical and ecological defense functions, spawning a new field of study called ecological biochemistry [KUHLSCH, POHNERT 2015; NELSON, WHITEHEAD 2021; SHEAR 2015; STANLEY *et al.* 2016]. PSMs provide a variety of functions, including disease, insect, and herbivore defense, stress response, and modulating organismal relationships [BARAK 2022]. Plant microbiomes, on the other hand, regulate plant metabolism and so play a role in many of the above-mentioned activities. Plants may influence their microbiome by secreting different metabolites, according to studies, and the microbiome, in turn, can alter the host plant's metabolome. However, little is understood about the interactions that influence phenotypic alterations [SINGH *et al.* 2019]. Phenolics, flavonoids, and alkaloids are a few of the secondary metabolites of plants that have been covered in this study.

PLANT DEFENSE THROUGH PSMs

Flavonoids and phenylpropanoids, which have distinct modes of action, are broadly dispersed chemicals in the plant defense system [HILDRETH *et al.* 2022]. Lots of antifungal agents address only six processes, the majority of which function in tandem with cell signaling molecules and have physiological effects or target

pathogen components such as reproductive system, DNA alkylation, and enzyme inhibition, etc. [LE ROY *et al.* 2016]. The majority of these compounds include phenolics with hydroxyl groups that are prone to disintegrate into phenolate ions [HADACEK 2002]. Because phenolic hydroxyl groups establish hydrogen and ionic connections with protons and peptides, the larger their number, the more denaturing and astringent the product [WINK 2018]. Proteins cannot function effectively without confirmation, i.e., a good three-dimensional structure. Any alteration in protein conformation alters the characteristics of the protein, preventing interaction with other RNA/DNA and proteins [MARSAFARI *et al.* 2020]. Secondary metabolites engage with free amino-, OH-, or SH- groups to form a covalent link, which changes the 3D structure of proteins, resulting in a change in protein turnover or loss of function [TIWARI *et al.* 2016]. Stronger ionic bonds and hydrogen bonds are formed by polyphenols [HAN *et al.* 2020]. The protein's flexibility is altered whenever these weak non-covalent bonds form at the same time and interact with a protein, resulting in protein inactivation. Since phenols' polar nature reduces their absorption after oral ingestion, they are less hazardous compounds [WANG *et al.* 2019]. A wood degradation test using stilbene impregnated birch and aspen samples was used to investigate pinosylvin's influence and resveratrol fungal growth [SEPPÄNEN *et al.* 2004]. Resveratrol stimulates the development of fungal growth at pinosylvin concentrations that are sufficient to prohibit fungal growth [GOYAL *et al.* 2012; SALEHI *et al.* 2018].

PSMs AS ANTIVIRAL, ANTIFUNGAL, AND ANTIBACTERIAL COMPOUNDS

Flavonoids, phenolics, and alkaloids, among other secondary plant metabolic products have antiviral properties [HUSSEIN, EL-ANSSARY 2018]. Alkaloids have various physiologically active molecules that affect living beings due to their diverse architectures. Antiviral activities have been discovered in over 18,000 alkaloids found in traditional Chinese herbs [JUBAIR *et al.* 2021]. The plantain lily (*Hosta plantaginea*) alkaloid 7-deoxy-trans-dihydronarciclasine (named E144) has an anti-TMV activity with a minimum IC₅₀ value of 1.80 μM [YANG, HE 2021]. Furthermore, Bruceine-D, which may be found in the Brassica javanica extract, has an inhibiting effect against TMV, CMV, and PVY [MAO *et al.* 2019]. White goosefoot (*Chenopodium amaranticolor*) extract embedded with 100 $\mu\text{g}\cdot\text{dm}^{-3}$ Bruceine-D has been shown to suppress more than 90% of CMV and PVY infection after 15 minutes of treatment [ALIEV, KURBANOVA 2022].

According to research, the majority of secondary metabolites are thought to have antifungal properties [BASILE *et al.* 2015]. Allied flavonoids and phenolics make up a wide set of phytochemicals [VELU *et al.* 2018]. As indicated in Table 1, these chemicals are found in large quantities in the leaves and skins of fruits and play a role in plant disease resistance, UV resistance, and defense against pigmentation.

Phenolics have been shown to alter microbe cell permeability and induce functional and structural distortion of membrane proteins, causing membrane bonded enzymes, ATP generation and preservation system, pH gradient, and substrate consumption for ATP synthesis to be disrupted [BENHAMOU *et al.* 1996]. The pathogens' development in the apoplast is slowed by

Table 1. List the antifungal properties of phenolic compounds

Fungus	Chemical compound
<i>Phytophthora infestans</i>	vanillic acid
<i>Phytophthora</i>	oleuropein
<i>Phoma tracheiphila</i>	nobiletin
<i>Penicillium digitatum</i>	naringin
<i>Monilinia fructicola</i>	genistein
<i>Helminthosporium sativum</i>	hordatol A
<i>Fusarium oxysporum</i>	chlorogenic acid
<i>Eutypa lata</i>	salicylic acid
<i>Colletotrichum circinans</i>	protocatechuic acid
<i>Botrytis cinerea</i>	benzaldehyde
<i>Aspergillus</i>	flavones

Source: own elaboration based on literature.

antimicrobial substances [Kwon et al. 2008]. Secondary metabolites may have antibacterial properties, according to one research [Helmann et al. 2022]. Legume leaf extracts in liquid form, such as *Zapoteca portoricensis*, were also shown to contain flavonoids. The antimicrobial substances' defensive role in the study of *Arabidopsis* root exudates infected with *Pseudomonas syringae* was identified in the root exudates as a defense mechanism against this bacterium [Helmann et al. 2022]. It was discovered that plants infected with strains that are not pathogenic produce a greater quantity of secondary metabolites than plants inoculated with virulent strains and that *P. syringae* is unable to infect *Arabidopsis* in seven out of eight strains [O'Malley, Anderson 2021; Valette et al. 2020]. Non-pathogenic bacteria have a mild antibacterial effect on non-infecting strains. Infecting strains had no effect on root exudation's antibacterial efficacy [Guo et al. 2021].

CONCLUSIONS

Due to their significant role in the physiology of plant stress, secondary metabolites have attracted a lot of attention. Even in challenging circumstances, these metabolites have the potential to preserve the health and productivity of food crops with little loss. By increasing the synthesis of numerous enzymes involved in the production of secondary metabolites and raising the expression of various genes involved in plant resistance mechanisms, PSMs mediate a variety of defensive functions. In this work, in order to control bacterial, viral, fungal, and insect diseases, we have highlighted the most recent developments in the use of secondary metabolites. With the passage of time, plants acquired defense systems against a variety of abiotic and biotic stressors. Apart from secondary metabolites that are either shown to be produced or caused by infection, sophisticated techniques are needed to accurately examine the relationship between crop resistance management and nitrogen and sulphur treatment. Previous research has shown that the quantity of certain substances in the growing medium affects the formation of N- and S-containing secondary metabolites, implying that an optimal amount of these nutrients is necessary for healthy plant development and resistance to environmental and microorganisms' challenges.

REFERENCES

- ALIEV R.I., KURBANOVA M.N. 2022. Prospects for the development of biotechnology: New directions, distribution and application. AIP Conference Proceedings. Vol. 2390(1), 030001. DOI 10.1063/5.0075804.
- ANSARI I.A., AKHTAR M.S. 2019. Current insights on the role of terpenoids as anticancer agents: A perspective on cancer prevention and treatment. In: Natural bio-active compounds. Vol. 2. Chemistry, pharmacology and health care practices. Eds. M.K. Swamy, M.S. Akhtar. Singapore. Springer p. 53–80. DOI 10.1007/978-981-13-7205-6_3.
- BADRI D.V., CHAPARRO J.M., ZHANG R., SHEN Q., VIVANCO J.M. 2013. Application of natural blends of phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. Journal of Biological Chemistry. Vol. 288(7) p. 4502–4512. DOI 10.1074/jbc.M112.433300.
- BARAK D. 2022. Role of plant metabolites in plant protection and their potential in integrated pest management. The Pharma Innovation. Vol. 11(5) p. 699–704. DOI 10.22271/tpi.2022.v11.i5i.12469.
- BARTWAL A., MALL R., LOHANI P., GURU S.K., ARORA S. 2013. Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. Journal of Plant Growth Regulation. Vol. 32(1) p. 216–232. DOI 10.1007/s00344-012-9272-x.
- BASILE A., RIGANO D., LOPPI S., DI SANTI A., NEBBIOSO A., SORBO S., ..., BONTEMPO P. 2015. Antiproliferative, antibacterial and antifungal activity of the lichen *Xanthoria parietina* and its secondary metabolite parietin. International Journal of Molecular Sciences. Vol. 16(4) p. 7861–7875. DOI 10.3390/ijms16047861.
- BELAIR M., GRAU A.L., CHONG J., TIAN X., LUO J., GUAN X., PENSEC F. 2022. Pathogenicity factors of botryosphaeriaceae associated with grapevine trunk diseases: New developments on their action on grapevine defense responses. Pathogens. Vol. 11(8), 951. DOI 10.3390/pathogens11080951.
- BENHAMOU N., KLOEPPER J.W., QUADT-HALLMAN A., TUZUN S. 1996. Induction of defense-related ultrastructural modifications in pea root tissues inoculated with endophytic bacteria. Plant Physiology. Vol. 112(3) p. 919–929. DOI 10.1104/pp.112.3.919.
- BLAŽEVIĆ I., MONTAUT S., BURČUL F., OLSEN C.E., BUROW M., ROLLIN P., AGERBIK N. 2020. Glucosinolate structural diversity, identification, chemical synthesis and metabolism in plants. Phytochemistry. Vol. 169, 112100. DOI 10.1016/j.phytochem.2019.112100.
- BODENHAUSEN N., REYMOND P. 2007. Signaling pathways controlling induced resistance to insect herbivores in *Arabidopsis*. Molecular Plant-Microbe Interactions®. Vol. 20(11) p. 1406–1420. DOI 10.1094/MPMI-20-11-1406.
- BRIBI N. 2018. Pharmacological activity of alkaloids: A review. Asian Journal of Botany. Vol. 1 p. 1–6.
- CHAKRABORTY N., ACHARYA K. 2017. “NO way”! Says the plant to abiotic stress. Plant Gene. Vol. 11 p. 99–105. DOI 10.1016/j.plgene.2017.05.001.
- CHEYNIER V., COMTE G., DAVIES K.M., LATTANZIO V., MARTENS S. 2013. Plant phenolics: Recent advances on their biosynthesis, genetics, and ecophysiology. Plant Physiology and Biochemistry. Vol. 72 p. 1–20. DOI 10.1016/j.plaphy.2013.05.009.
- DIVEKAR P.A., NARAYANA S., DIVEKAR B.A., KUMAR R., GADRATAGI B.G., RAY A., ..., BEHERA T.K. 2022. Plant secondary metabolites as defense tools against herbivores for sustainable crop protection. International Journal of Molecular Sciences. Vol. 23(5), 2690. DOI 10.3390/ijms23052690.

- DIXON D.P., SKIPSEY M., EDWARDS R. 2010. Roles for glutathione transferases in plant secondary metabolism. *Phytochemistry*. Vol. 71(4) p. 338–350. DOI 10.1016/j.phytochem.2009.12.012.
- GALCZYŃSKA M., MAŃKOWSKA N., MILKE J., BUŚKO M. 2019. Possibilities and limitations of using *Lemna minor*, *Hydrocharis morsus-ranae* and *Ceratophyllum demersum* in removing metals with contaminated water. *Journal of Water and Land Development*. No. 40. DOI 10.2478/jwld-2019-0018.
- GALILI G., AMIR R., HOEFGEN R., HESSE H. 2005. Improving the levels of essential amino acids and sulfur metabolites in plants. *Biological Chemistry*. Vol. 386(9) p. 817–831. DOI 10.1515/BC.2005.097.
- GARCIA-MIER L., JIMENEZ-GARCÍA S.N., SALAZAR C.S., CONTRERAS-MEDINA L.M., ESCALANTE K.E., MARTINEZ C.G., GARCÍA-TREJO J.F., GUEVARA-GONZALEZ R.G., FERERINO-PEREZ A.A. 2019. Strategies that influence the production of secondary metabolites in plants. In: *Nutritional quality improvement in plants concepts and strategies in plant sciences*. Eds. P.K. Jaiwal, A.K. Chhillar, D. Chaudhary, R. Jaiwal p. 231–270. Cham. Springer International Publishing. DOI 10.1007/978-3-319-95354-0_9.
- GOYAL S., LAMBERT C., CLUZET S., MÉRILLON J.M., RAMAWAT K.G. 2012. Secondary metabolites and plant defence. In: *Plant defence: Biological control progress in biological control*. Eds. J.M. Mérillon, K.G. Ramawat p. 109–138. Dordrecht. Springer Netherlands. DOI 10.1007/978-94-007-1933-0_5.
- GUO X., WAN Y., SHAKEEL M., WANG D., XIAO L. 2021. Effect of mycorrhizal fungi inoculation on bacterial diversity, community structure and fruit yield of blueberry. *Rhizosphere*. Vol. 19, 100360. DOI 10.1016/j.rhisph.2021.100360.
- HADACEK F. 2002. Secondary metabolites as plant traits: Current assessment and future perspectives. *Critical Reviews in Plant Sciences*. Vol. 21(4) p. 273–322. DOI 10.1080/0735-260291044269.
- HAMMER T.J., BOWERS M.D. 2015. Gut microbes may facilitate insect herbivory of chemically defended plants. *Oecologia*. Vol. 179(1) p. 1–14. DOI 10.1007/s00442-015-3327-1.
- HAN Y., LIN Z., ZHOU J., YUN G., GUO R., RICHARDSON J.J., CARUSO F. 2020. Polyphenol-mediated assembly of proteins for engineering functional materials. *Angewandte Chemie International Edition*. Vol. 59(36) p. 15618–15625. DOI 10.1002/anie.202002089.
- HEDJAL S., ZOUINI D., BENAMARA A. 2018. Hydrochemical assessment of water quality for irrigation: A case study of the wetland complex of Guerbes-Sanhadja, North-East of Algeria. *Journal of Water and Land Development*. No. 38. DOI 10.2478/jwld-2018-0041.
- HELMANN T.C., KING D.M., LINDOW S.E. 2022. Differential virulence contributions of the efflux transporter MexAB-OprM in *Pseudomonas syringae* infecting a variety of host plants. *Molecular Plant-Microbe Interactions*. Vol. 35(8) p. 672–680. DOI 10.1094/MPMI-04-21-0099-R.
- HILDRETH S.B., LITTLETON E.S., CLARK L.C., PULLER G.C., KOJIMA S., WINKEL B.S.J. 2022. Mutations that alter *Arabidopsis* flavonoid metabolism affect the circadian clock. *The Plant Journal*. Vol. 110(4) p. 932–945. DOI 10.1111/tpj.15718.
- HILLIER S.G., LATHE R. 2019. Terpenes, hormones and life: Isoprene rule revisited. *Journal of Endocrinology*. Vol. 242(2) p. R9–R22. DOI 10.1530/JOE-19-0084.
- HUSSEIN R.A., EL-ANSSARY A.A. 2018. Plants secondary metabolites: The key drivers of the pharmacological actions of medicinal plants. In: *Herbal medicine*. Ed. P.F. Builders. IntechOpen p. 11–30. DOI 10.5772/intechopen.76139.
- JAMLOKI A., BHATTACHARYA M., NAUTIYAL M.C., PATNI B. 2021. Elucidating the relevance of high temperature and elevated CO₂ in plant secondary metabolites (PSMs) production. *Heliyon*. Vol. 7(8), e07709. DOI 10.1016/j.heliyon.2021.e07709.
- JUBAIR N., RAJAGOPAL M., CHINNAPPAN S., ABDULLAH N.B., FATIMA A. 2021. Review on the antibacterial mechanism of plant-derived compounds against multidrug-resistant bacteria (MDR). *Evidence-Based Complementary and Alternative Medicine*. Vol. 2021, e3663315. DOI 10.1155/2021/3663315.
- KEITH R.A., MITCHELL-OLDS T. 2017. Testing the optimal defense hypothesis in nature: Variation for glucosinolate profiles within plants. *PLOS ONE*. Vol. 12(7), e0180971. DOI 10.1371/journal.pone.0180971.
- KILGORE M.B., KUTCHAN T.M. 2016. The *Amaryllidaceae* alkaloids: Biosynthesis and methods for enzyme discovery. *Phytochemistry Reviews*. Vol. 15(3) p. 317–337. DOI 10.1007/s11101-015-9451-z.
- KOZA N.A., ADEDAYO A.A., BABALOLA O.O., KAPPO A.P. 2022. Microorganisms in plant growth and development: Roles in abiotic stress tolerance and secondary metabolites secretion. *Microorganisms*. Vol. 10(8), 1528. DOI 10.3390/microorganisms10081528.
- KUHLISCH C., POHNERT G. 2015. Metabolomics in chemical ecology. *Natural Product Reports*. Vol. 32(7) p. 937–955. DOI 10.1039/C5NP00003C.
- KWON C., BEDNAREK P., SCHULZE-LEFERT P. 2008. Secretory pathways in plant immune responses. *Plant Physiology*. Vol. 147(4) p. 1575–1583. DOI 10.1104/pp.108.121566.
- LE ROY J., HUSS B., CREACH A., HAWKINS S., NEUTELINGS G. 2016. Glycosylation is a major regulator of phenylpropanoid availability and biological activity in plants. *Frontiers in Plant Science*. Vol. 7, 735. DOI 10.3389/fpls.2016.00735.
- LI Y., KONG D., FU Y., SUSSMAN M.R., WU H. 2020. The effect of developmental and environmental factors on secondary metabolites in medicinal plants. *Plant Physiology and Biochemistry*. Vol. 148 p. 80–89. DOI 10.1016/j.plaphy.2020.01.006.
- LICHMAN B.R. 2021. The scaffold-forming steps of plant alkaloid biosynthesis. *Natural Product Reports*. Vol. 38(1) p. 103–129. DOI 10.1039/D0NP00031K.
- MANNIELLO M.D., MORETTA A., SALVIA R., SCIEUZO C., LUCCHETTI D., VOGEL H., SGAMBATO A., FALABELLA P. 2021. Insect antimicrobial peptides: Potential weapons to counteract the antibiotic resistance. *Cellular and Molecular Life Sciences*. Vol. 78(9) p. 4259–4282. DOI 10.1007/s00018-021-03784-z.
- MAO G., TIAN Y., SUN Z., OU J., XU H. 2019. Bruceine D isolated from *Brucea javanica* (L.) Merr. as a systemic feeding deterrent for three major lepidopteran pests. *Journal of Agricultural and Food Chemistry*. Vol. 67(15) p. 4232–4239. DOI 10.1021/acs.jafc.8b06511.
- MARSAFARI M., SAMIZADEH H., RABIEI B., MEHRABI A., KOFFAS M., XU P. 2020. Biotechnological production of flavonoids: An update on plant metabolic engineering, microbial host selection, and genetically encoded biosensors. *Biotechnology Journal*. Vol. 15(8), 1900432. DOI 10.1002/biot.201900432.
- MICHAEL J.P. 2016. Chapter one – Simple indolizidine and quinolizidine alkaloids. In: *The alkaloids: Chemistry and biology*. Vol. 75. Ed. H.-J. Knölker. Academic Press. ISBN 9780128034347 pp. 498. DOI 10.1016/bs.alkal.2014.12.001.
- NELSON A.S., WHITEHEAD S.R. 2021. Fruit secondary metabolites shape seed dispersal effectiveness. *Trends in Ecology & Evolution*. Vol. 36(12) p. 1113–1123. DOI 10.1016/j.tree.2021.08.005.
- NINKUU V., ZHANG L., YAN J., FU Z., YANG T., ZENG H. 2021. Biochemistry of terpenes and recent advances in plant protection. *International Journal of Molecular Sciences*. Vol. 22(11), 5710. DOI 10.3390/ijms22115710.
- O'MALLEY M.R., ANDERSON J.C. 2021. Regulation of the *Pseudomonas syringae* type III secretion system by host environment signals.

- Microorganisms. Vol. 9(6), 1227. DOI 10.3390/microorganisms9061227.
- PANDEY A., PATHAK J., SINGH D.K., AHMED H., SINGH V., KUMAR D., SINHA R.P. 2020. Photoprotective role of UV-screening pigment scytonemin against UV-B-induced damages in the heterocyst-forming cyanobacterium *Nostoc* sp. strain HKAR-2. Brazilian Journal of Botany. Vol. 43(1) p. 67–80. DOI 10.1007/s40415-020-00589-5.
- PIASECKA A., JEDRZEJCZAK-REY N., BEDNAREK P. 2015. Secondary metabolites in plant innate immunity: Conserved function of divergent chemicals. New Phytologist. Vol. 206(3) p. 948–964. DOI 10.1111/nph.13325.
- PRASANTH C.N., VISWANATHAN R., MALATHI P., SUNDAR A.R. 2022. Carbohydrate active enzymes (CAZy) regulate cellulolytic and pectinolytic enzymes in *Colletotrichum falcatum* causing red rot in sugarcane. 3 Biotech. Vol. 12(2), 48. DOI 10.1007/s13205-022-03113-6.
- RAJPUT A., SHARMA R., BHARTI R. 2022. Pharmacological activities and toxicities of alkaloids on human health. Materials Today: Proceedings. Vol. 48 p. 1407–1415. DOI 10.1016/j.matpr. 2021.09.189.
- RODRIGUES-CORRÊA K.C. DA S., FETT-NETO A.G. 2019. Abiotic stresses and non-protein amino acids in plants. Critical Reviews in Plant Sciences. Vol. 38(5–6) p. 411–430. DOI 10.1080/07352689.2019.1707944.
- SALEHI B., MISHRA A.P., NIGAM M., SENER B., KILIC M., SHARIFI-RAD M., FOKOU P.V.T., MARTINS N., SHARIFI-RAD J. 2018. Resveratrol: A double-edged sword in health benefits. Biomedicines. Vol. 6 (3), 91. DOI 10.3390/biomedicines6030091.
- SEPPÄNEN S.-K., SYRJÄLÄ L., VON WEISENBERG K., TEERI T.H., PAAJANEN L., PAPPINEN A. 2004. Antifungal activity of stilbenes in in vitro bioassays and in transgenic *Populus* expressing a gene encoding pinosylvin synthase. Plant Cell Reports. Vol. 22(8) p. 584–593. DOI 10.1007/s00299-003-0728-0.
- SHARMA A., SHARMA S., KUMAR A., KUMAR V., SHARMA A.K. 2022. Plant secondary metabolites: An introduction of their chemistry and biological significance with physicochemical aspect. In: Plant secondary metabolites: Physico-chemical properties and therapeutic applications. Eds. A.K. Sharma, A. Sharma p. 1–45. Singapore. Springer Nature. DOI 10.1007/978-981-16-4779-6_1.
- SHEAR W.A. 2015. The chemical defenses of millipedes (diplopoda): Biochemistry, physiology and ecology. Biochemical Systematics and Ecology. Vol. 61 p. 78–117. DOI 10.1016/j.bse.2015.04.033.
- SINGH D., RAINA T.K., KUMAR A., SINGH J., PRASAD R. 2019. Plant microbiome: A reservoir of novel genes and metabolites. Plant Gene. Vol. 18, 100177. DOI 10.1016/j.plgene.2019.100177.
- STANLEY C.E., GROSSMANN G., SOLVAS X.C. I., DEMELLO A.J. 2016. Soil-on-a-Chip: microfluidic platforms for environmental organismal studies. Lab on a Chip. Vol. 16(2) p. 228–241. DOI 10.1039/C5LC01285F.
- TIWARI P., SANGWAN R.S., SANGWAN N.S. 2016. Plant secondary metabolism linked glycosyltransferases: An update on expanding knowledge and scopes. Biotechnology Advances. Vol. 34(5) p. 714–739. DOI 10.1016/j.biotechadv.2016.03.006.
- TSUDA K., KATAGIRI F. 2010. Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. Current Opinion in Plant Biology. Vol. 13(4) p. 459–465. DOI 10.1016/j.pbi.2010.04.006.
- TSUNODA T., KROSSE S., VAN DAM N.M. 2017. Root and shoot glucosinolate allocation patterns follow optimal defence allocation theory. Journal of Ecology. Vol. 105(5) p. 1256–1266. DOI 10.1111/1365-2745.12793.
- TYSKIEWICZ R., NOWAK A., OZIMEK E., JAROSZUK-ŚCISEL J. 2022. Trichoderma: The current status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. International Journal of Molecular Sciences. Vol. 23(4), 2329. DOI 10.3390/ijms23042329.
- UL HAQ S., KHAN A., ALI M., KHATTAK A.M., GAI W.-X., ZHANG H.-X., WEI A.-M., GONG Z.-H. 2019. Heat shock proteins: Dynamic biomolecules to counter plant biotic and abiotic stresses. International Journal of Molecular Sciences. Vol. 20(21), 5321. DOI 10.3390/ijms20215321.
- VALETTE M., REY M., GERIN F., COMTE G., WISNIEWSKI-DYÉ F. 2020. A common metabolomic signature is observed upon inoculation of rice roots with various rhizobacteria. Journal of Integrative Plant Biology. Vol. 62(2) p. 228–246. DOI 10.1111/jipb.12810.
- VELU G., PALANICHAMY V., RAJAN A.P. 2018. Phytochemical and pharmacological importance of plant secondary metabolites in modern medicine. In: Bioorganic phase in natural food: An overview. Eds. S.M. Roopan, G. Madhumitha p. 135–156. Cham. Springer International Publishing. DOI 10.1007/978-3-319-74210-6_8.
- VERMA N., SHUKLA S. 2015. Impact of various factors responsible for fluctuation in plant secondary metabolites. Journal of Applied Research on Medicinal and Aromatic Plants. Vol. 2(4) p. 105–113. DOI 10.1016/j.jarmap.2015.09.002.
- WAHEED A., HAXIM Y., ISLAM W., KAHAR G., LIU X., ZHANG D. 2022. Role of pathogen's effectors in understanding host-pathogen interaction. Biochimica et Biophysica Acta (BBA) – Molecular Cell Research. Vol. 1869(12), 119347. DOI 10.1016/j.bbamcr. 2022.119347.
- WANG Z., AN G., ZHU Y., LIU X., CHEN Y., WU H., WANG Y., SHI X., MAO C. 2019. 3D-printable self-healing and mechanically reinforced hydrogels with host-guest non-covalent interactions integrated into covalently linked networks. Materials Horizons. Vol. 6(4) p. 733–742. DOI 10.1039/C8MH01208C.
- WAR A.R., BUHROO A.A., HUSSAIN B., AHMAD T., NAIR R.M., SHARMA H. C. 2020. Plant defense and insect adaptation with reference to secondary metabolites. In: Co-evolution of secondary metabolites reference series in phytochemistry. Eds. J.-M. Mérillon, K.G. Ramawat Cham. Springer International Publishing p. 795–822. DOI 10.1007/978-3-319-96397-6_60.
- WINK M. 2018. Plant secondary metabolites modulate insect behavior-steps toward addiction? Frontiers in Physiology. Vol. 9, 364. DOI 10.3389/fphys.2018.00364.
- WU S., SHAN L., HE P. 2014. Microbial signature-triggered plant defense responses and early signaling mechanisms. Plant Science. Vol. 228 p. 118–126. DOI 10.1016/j.plantsci.2014.03.001.
- YANG L., HE J. 2021. Traditional uses, phytochemistry, pharmacology and toxicological aspects of the genus *Hosta* (Liliaceae): A comprehensive review. Journal of Ethnopharmacology. Vol. 265, 113323. DOI 10.1016/j.jep.2020.113323.
- YESHI K., CRAYN D., RITMEJERYTÉ E., WANGCHUK P. 2022. Plant secondary metabolites produced in response to abiotic stresses has potential application in pharmaceutical product development. Molecules. Vol. 27(1) p. 313. DOI 10.3390/molecules27010313.
- ZHANG Y., DENG T., SUN L., LANDIS J.B., MOORE M.J., WANG H., ..., SUN H. 2021. Phylogenetic patterns suggest frequent multiple origins of secondary metabolites across the seed-plant 'tree of life'. National Science Review. Vol. 8(4), nwaa105. DOI 10.1093/nsr/nwaa105.